

## Influence of elevated CO<sub>2</sub> concentrations on the growth of *Armillaria ostoyae* (Romagn.) Herink rhizomorphs *in vitro*

Paweł Lech<sup>1\*</sup>, Anna Żółciak<sup>2</sup> 

Forest Research Institute, <sup>1</sup>Department of Forest Resources Management,  
<sup>2</sup>Department of Forest Protection, Sękocin Stary, ul. Braci Leśnej 3, 05–090 Raszyn, Poland

\*Tel. +48 22 7153825, fax +48 22 7153837, e-mail: P.Lech@ibles.waw.pl

**Abstract.** A comparative experiment was carried out in growth chambers to determine the effects of elevated CO<sub>2</sub> concentrations (either 760 ppm or 1,140 ppm) versus ambient CO<sub>2</sub> conditions on the growth of *Armillaria ostoyae* (Romagn.) Herink rhizomorphs, which is the infectious organ of a fungal pathogen affecting many forest trees. We found that one out of three isolates in the experiment differed significantly in rhizomorph production, which was measured as rhizomorph dry mass/100 days of growth. Rhizomorph production was also affected by the tree species used as a food source in the inoculum preparation, with beech wood being significantly different from oak and hazel. Under higher CO<sub>2</sub> regimes the production of rhizomorphs was consistently lower for all three isolates compared to ambient CO<sub>2</sub> concentrations. For one isolate (no. 11) the growth differences were significant between 380 ppm and both elevated CO<sub>2</sub> concentrations (760 ppm and 1,140 ppm), while for the other two (no. 30 and 32) significance was observed only between 380 ppm and 760 ppm. No statistically significant differences have been noted between 760 ppm and 1,140 ppm CO<sub>2</sub> for these two isolates. It was concluded that elevated concentrations of CO<sub>2</sub> inhibited *A. ostoyae* rhizomorph growth and therefore have the potential to lessen the pathogenicity of the fungus.

**Keywords:** *Armillaria ostoyae*, CO<sub>2</sub>, rhizomorph growth

### 1. Introduction

Rhizomorphs are infectious organs of honey fungi, including the most destructive pathogen of this genus affecting most of forest tree species in Poland – *Armillaria ostoyae* (Romagn.) Herink. Next to the contacts of mycelium and spores with tree roots, rhizomorphs add up to propagation of the so-called ‘white rot’ root disease caused by honey fungi in forest stands (Żółciak 1999, 2005). For this reason, the morphology and functions of rhizomorphs, as well as rhizomorphogenesis, have long been the subject of numerous studies (Hartig 1873; Manka 1953; Jacques-Felix 1967, 1968; Redfern 1973; Rykowski 1984; Redfern, Filip 1991; Łakomy 2004; Guillaumin, Botton 2005; Guillaumin, Legrand 2005; Lung-Escarmant et al. 2005). Amongst other things, the influence of several substances, such as ethanol and other alcohols, growth substances (o-aminobenzoic acid, p-palminobenzoic acid, indole-3-acetic acid) (Pentland

1965, 1967; Garraway, Weinhold 1967, 1968 a, b; Garraway 1970; Sortkjaer, Allermann 1972a, b, c) and also inorganic nutrients (Morrison 1975, Rykowski 1984, Przybył 1998), on the formation and growth of rhizomorph has been studied. Furthermore, the effects of soil moisture and temperature on the rhizomorphogenesis have been demonstrated. Appropriate humidity conditions boost proper functioning of the apical meristem and thus stimulate the growth of rhizomorph (Rykowski 1984). The temperature affects not only the initiation and development but also the number and extent of branching of rhizomorph (Redfern 1973). The response of rhizomorphs to the presence of oxygen and carbon dioxide in the soil environment was also reported (Hintikka, Korhonen 1970; Hintikka 1974; Morrison 1976). According to Morrison (1976), rhizomorphs prefer the anaerobic environment and the direction of their growth in a given soil profile depends on fluctuations in oxygen and carbon dioxide pressure. Hintikka (1974) showed the stimulatory effect

Submitted: 2.02.2017, reviewed: 29.03.2017, accepted after revision: 26.04.2017

of CO<sub>2</sub> on the growth of rhizomorphs, which was explained by their expansion towards the roots that diffuse CO<sub>2</sub> into the soil. Olszyk et al. (2001) observed initial fungal colonies in forest soil and noted their enhanced formation and growth under the influence of elevated concentrations of atmospheric CO<sub>2</sub>. The ability of wood-decay fungi, including honey fungi, to assimilate CO<sub>2</sub> was also experimentally demonstrated; however, the assimilation intensity was low: 1 g of the fungal tissue structure contained about 0.017 mg of assimilated carbon (Schinner, Concin 1981).

In the coming decades, the increase in atmospheric CO<sub>2</sub> (already observed in the past 200 years) will go on faster and faster to reach around 490–1,250 ppm at the end of this century – depending on the adopted variant of fossil fuel consumption and implemented models of economic development (Climate Change 2001). The study results concerning the effects of elevated CO<sub>2</sub> on the rhizomorphogenesis and rhizomorph production of *A. ostoyae*, as well as those regarding the infection potential of this fungus, have been hardly available. The results obtained by Lech and Żółciak (2006, 2017) have been published – however, these seem not explicit enough. The results of the first study (2006) showed the growth stimulation of rhizomorphs under elevated CO<sub>2</sub> concentration (750 ppm), but pine seedlings were not yet infected or killed by the fungus examined. The results of the second study (2017) showed that the increase in CO<sub>2</sub> concentration to 1,000 ppm slowed down the disease process, as well as the mortality of 2-year-old pine seedlings and 3-year-old spruce seedlings was decreased 16 months after artificial inoculation with *A. ostoyae*.

Not too many experimental studies have been so far devoted to the impacts of climate change on other fungal pathogens of forest trees. Those that have been undertaken were focused on the effect of increased concentrations of CO<sub>2</sub> on threat patterns in oak and beech populations due to *Phytophthora* species (Fleischman et al. 2010; Tkaczyk et al. 2014; Oszako et al. 2016), as well as those in poplar populations due to *Melampsora medusa* Thümen – the causal agent of poplar rust (Percy et al. 2002). Then again, the results obtained were inconclusive, as both the disease stimulation as well as no effects of elevated CO<sub>2</sub> on the course of the disease process were observed.

Scarcity of studies with regard to the impact of predicted climate change on forest fungal tree pathogens, as well as incompleteness and ambiguity of the results obtained, have been noted in subject literature reviews (Manning, von Tiedemann 1995; Chakraborty et al. 2000; Garrett et al. 2006). The authors emphasise that further research is the only way to fill knowledge gaps in this field. This argument was the reason behind undertaking the present study. The null hypothesis assumed that elevated CO<sub>2</sub> would have stimulating effect on the growth of *A. ostoyae* rhizomorphs.

## 2. Material and Methods

In order to verify the null hypothesis, the experiment was carried out under artificial, controlled environment. The development of *A. ostoyae* rhizomorphs was initiated using inoculum placed in glass cylindrical containers (opened at both ends). These were filled with medium-sized grains quartz sand and sterilised before the placement of the inoculum inside. The containers with inoculum were placed in the climatic chambers and exposed to the atmosphere with the following CO<sub>2</sub> concentrations for the next 6–7 months: 380 (control, nearly ambient CO<sub>2</sub>), 760 and 1,140 ppm (2- and 3-fold of control CO<sub>2</sub> concentration, respectively). Other than that, the conditions for all the experimental variants were maintained the same for the duration of the experiment, that is, constant relative humidity (80%) and temperature (22°C). In order to ensure the adequate moisture of inoculum substrate (sand), irrigation with deionised water was applied twice a week. The experiment was conducted with the use of the climatic chambers Mytron WB 750.

Three *A. ostoyae* isolates were used for inoculum production. The isolates were obtained from 3 different spruce stands, growing in the Forest Districts: Ujsoly located in the Żywiec Beskid Mts. and Wisła – in the Silesian Beskid Mts. The fungal material for research was collected and the species was classified based on the morphological features of fruiting bodies (Żółciak 1999) as well as pure cultures were identified by means of intersterility tests (Korhonen, 1978). Beech, oak and hazel wood were used as food base for *A. ostoyae*. The inoculum was prepared consistent with the methodology by Rykowski (1984) and Redfern and Filip (1991). Tree branches, 5–7 cm long and 1.5–2 cm in diameter, were inoculated with *A. ostoyae* isolates, after washing and autoclaving for 40 minutes at 108°C under 0.05-MPa pressure and re-sterilizing on the next day (following the above procedure). The incubation lasted approximately 3 months.

The dry mass of rhizomorphs was determined using Sartorius analytic A200S balance (accurateness 0.0001 g) approximately 6–7 months after placing the containers with the inocula into the climatic chambers and was adopted as criterion of rhizomorph development. In order to perform the comparisons and statistical analyses, dry mass results were expressed per 100 days of rhizomorph growth. Owing to the large number of observations and rhizomorph brittleness, evaluations of other biometric parameters, such as the length or the number of active peaks, were neglected.

In total, the study comprised 177 observations. The number of observations ranged from 28 to 86 for individual main factors, and that for interactions ranged from 10 to 30 (Table 1). The experiment might be described with the following model:

$$SM = \text{Isolate} + CO_2 + \text{Tree species used for inoculum production} + \text{interaction (Isolate} \times CO_2)$$

Where:

$SM$  is the rhizomorph dry mass per 100 days of growth,  
*Isolate* is the *A. ostoyae* isolate used for artificial inoculation,  
 $CO_2$  is the  $CO_2$  concentration tested,  
*Tree species used for inoculum production* is the wood of a given tree species used as inoculum food base.

The results obtained were statistically analysed by means of analysis of variance ANOVA with the main effects: *A. ostoyae* isolate (No. 11, 30 and 32),  $CO_2$  concentration (near-ambient, approximately 380 ppm; elevated, 760 and 1,140 ppm) and tree species used for inoculum production (beech, oak and hazel), as well as interaction effect (*A. ostoyae* isolate)  $\times$  ( $CO_2$  concentration). In order to meet ANOVA assumptions (normality of variable distribution and equality of variances for the tested variant combinations), data was subjected to logarithmic transformation. Tukey's highest significant difference HSD test was used to compare the means obtained in experimental variants (rhizomorph dry mass/100 days of growth), taking into account adjustments due to different numbers of observations in the variants of the experiment. The analyses were carried out using Statgraphics Centurion XV software.

### 3. Results

Statistical characteristics of the results with respect to rhizomorph dry matter/100 days of growth are presented in Table 1. Out of the three pathogen isolates used in the experiment, the largest dry mass of rhizomorphs was produced by the isolate No. 11 (0.02186 g), whereas other two isolates (No. 30 and 32) showed clearly lower values for this parameter (0.01229 and 0.00909 g, respectively). The greatest dry weight values were observed in rhizomorphs growing under control (near-ambient) concentration of  $CO_2$  (0.02923 g). At higher  $CO_2$  concentrations (760 and 1,140 ppm), rhizomorph dry mass was lower (0.00549 and 0.00852 g, respectively). All three isolates showed the highest dry mass of rhizomorphs/100 days of growth under 380 ppm of  $CO_2$ . At higher  $CO_2$  concentrations, rhizomorph dry mass values were lower, and the lowest dry mass values were observed in all the isolates growing under 760 ppm of  $CO_2$ . Under 380 ppm of  $CO_2$ , the isolate No. 11 showed the highest mean rhizomorph dry weight (0.05240 g), then the isolate No. 30 (0.02174 g; almost 2.5-fold lower as compared to No.11) and the isolate No. 32 (0.01355 g; nearly 4-fold lower as compared to No.11). Rhizomorphs growing under 760 ppm of  $CO_2$  showed the highest mean dry mass value in the case of the isolate No.30 (0.00740 g), then No. 32 (0.00558 g; 50% of No. 30), whereas dry mass of the isolate No. 11 was the lowest (0.00349 g). In the case of the highest  $CO_2$  concentration tested (1,140 ppm), the highest mean dry weight

of rhizomorphs was observed in the isolate No. 11 (0.00968 g); it was lower in the isolate No. 32 (0.00815 g) and the lowest in the isolate No. 30 (0.00774 g) (Table 1).

ANOVA results are presented in Figures 1 and 2 and Table 2. The analysis showed statistically significant differences in the production of *A. ostoyae* rhizomorphs per 100 days of growth, for all three main effects: *Isolate*, *Tree species used for the production of inoculum* and  $CO_2$ , as well as in the case of interaction *Isolate*  $\times$   $CO_2$  (Table 2). As shown by the results of Tukey's test, the isolate No. 11 as well as the rhizomorphs initiated from inoculum growing on beech wood showed significantly different means of rhizomorph dry weight. The growth of rhizomorphs significantly differed depending on the  $CO_2$  concentration, reaching the highest values under 380 ppm of  $CO_2$  and the lowest under 760 ppm of  $CO_2$ . The differences between dry mass values obtained for rhizomorphs of the isolates No. 30 and No. 32 were not statistically significant, as were the differences between the mean rhizomorph dry mass values observed when oak wood and hazel wood were used for inoculum production (Fig. 1).

With regard to the interaction *Isolate*  $\times$   $CO_2$ , Tukey's test showed the significance of differences between the means of rhizomorph dry mass/100 days of growth. For the isolate No. 11, these differences were significant between 380 ppm of  $CO_2$  and both elevated concentrations (760 and 1,140 ppm), whereas for the isolates No. 30 and No. 32, the significant differences were observed only between 380 and 760 ppm. In the case of the isolates No. 11 and No. 30, no significant differences in rhizomorph dry mass were found between 760 and 1,140 ppm of  $CO_2$ , and in the case of isolate No. 32, no significant differences in rhizomorph dry mass were found between 380 and 1,140 ppm (Figure 2).

### 4. Discussion


The results of the study and statistical analyzes carried out did not confirm the null hypothesis. Elevated  $CO_2$  concentrations (760 and 1,140 ppm) did not stimulate the growth of *A. ostoyae* rhizomorphs; on the contrary, rhizomorph growth was somewhat inhibited under these conditions. In all three pathogen isolates tested, the largest dry mass of rhizomorphs was recorded under near-ambient  $CO_2$  concentration (380 ppm). For one isolate, the differences between ambient and both elevated  $CO_2$  concentrations tested were statistically significant, and for the remaining two, the differences were significant between near-ambient  $CO_2$  and one of the elevated  $CO_2$  concentrations (760 ppm). These results correspond to those of the earlier study by Lech and Żółciak (2017), who showed the slowed-down course of the disease process as well as reduced mortality of pine and spruce seedlings

**Table 1.** Mean values of rhizomorphs dry mass per 100 days of growth and confidence limits for means for main effects (isolation of *A. ostoyae*, level of air CO<sub>2</sub> and tree species used for inoculum production) as well as for interaction: *isolate x level CO<sub>2</sub>*

Sources of variation	Number of observations	Mean SM / 100 days [g]	95% confidence intervals for means		
			lower	upper	
Experiment total	177	0.01441	0.01363	0.01988	
Isolate no.	11	86	0.02186	0.01906	0.02465
	30	44	0.01229	0.00777	0.01681
	32	47	0.00909	0.00480	0.01338
CO <sub>2</sub> [ppm]	380	66	0.02923	0.02568	0.03277
	760	52	0.00549	0.00137	0.00961
	1140	59	0.00852	0.00473	0.01232
Tree species used for inoculum production	Bk	28	0.01449	0.00879	0.02019
	Db	74	0.01430	0.01122	0.01738
	Lesz	75	0.01445	0.01143	0.01746
Interaction (Isolate <sup>x</sup> CO <sub>2</sub> )	11 <sup>x</sup> 380	30	0.05240	0.04767	0.05713
	11 <sup>x</sup> 760	27	0.00349	-0.00150	0.00847
	11 <sup>x</sup> 1140	29	0.00968	0.00487	0.01449
	30 <sup>x</sup> 380	19	0.02174	0.01547	0.02800
	30 <sup>x</sup> 760	10	0.00740	-0.00107	0.01587
	30 <sup>x</sup> 1140	15	0.00774	0.00076	0.01471
	32 <sup>x</sup> 380	17	0.01355	0.00695	0.02015
	32 <sup>x</sup> 760	15	0.00558	-0.00141	0.01257
	32 <sup>x</sup> 1140	15	0.00815	0.00117	0.01513


**Table 2.** Analysis of variance for rhizomorphs dry mass and adopted model of experiment. Statistically significant differences were marked with bold fonts.

Sources of variance	Sum of squares	Degree of freedom	Mean square	F	P
A: Isolate	46.51	2	23.25	15.19	<b>&lt;0.001</b>
B: CO <sub>2</sub>	134.42	2	67.21	43.90	<b>&lt;0.001</b>
C: Tree species used for inoculum production	13.83	2	6.91	4.52	<b>0.012</b>
Interaction (AB)	23.26	4	5.81	3.80	<b>0.006</b>
Residuals	254.15	166	1.53		
Total	505.02	176			



**Figure 1.** Means of rhizomorphs dry mass (after logarithmic transformation) and 95% confidence intervals for main effects of experiment: isolate of *A. ostoyae*, tree species of wood used for inoculum production and level of CO<sub>2</sub> used in the experiment

subjected to artificial inoculation with *A. ostoyae* and grown under elevated CO<sub>2</sub> (up to 1,000 ppm). In the latter study, as



**Figure 2.** Means of rhizomorphs dry mass (after logarithmic transformation) and 95% confidence intervals for interaction: isolates of *A. ostoyae* (no. 11, 30 and 32) and level of CO<sub>2</sub> used in the experiment (380 ppm, 760 ppm and 1,140 ppm)

a possible explanation of lower seedling mortality, the compensatory hypothesis was presented (Herms, Mattson 1992), assuming the targeted allocation of increased plant resources to actively defend against stress factors. This does not exclude, however, that in addition to the compensatory effect of increased CO<sub>2</sub> concentration on plants, raised CO<sub>2</sub> concentrations also inhibit rhizomorph growth and thus reduce the infection potential of the pathogen. Both of these phenomena may, therefore, contribute to slowing down the course of the disease process and decreasing seedling dieback.

The results of the study are fundamentally different from the results of another earlier study by Lech and Żółciak (2006), which showed a greater production of rhizomorphs from inoculum placed in the containers with pine seedlings grown under 750 ppm of CO<sub>2</sub>. Then, no contact rhizomorphs–tree roots and no infection of pine seedlings were observed; therefore, it was concluded that enhanced rhizomorph growth resulted from stimulating effect of raised CO<sub>2</sub> concentration. The ability of honey fungi to assimilate carbon dioxide (Schinner, Concini 1981) could support this explanation. At present, it seems that the reason behind rhizomorph growth enhancement could be higher air humidity in the climatic chambers with increased concentration of CO<sub>2</sub>. During the previous study (Lech, Żółciak 2006), the examined seedlings were grown in Mytron WB 750 chambers, using the B-647 controller (MKS Instruments) and a special hermetic container to maintain the assumed, increased concentration of CO<sub>2</sub>. The container was not equipped to regulate the relative humidity level, and this could have caused the increase/persistence of long-standing high moisture after watering the substrate, on which the seedlings were grown and *A. ostoyae* inoculum was

placed. The high moisture content of the substrate, as shown experimentally, is a factor favouring the growth of honey fungi rhizomorphs (Redfern 1973, Rykowski 1984).

## 5. Conclusions

The results of the study allow to draw the following conclusions:

- Raised CO<sub>2</sub> concentrations (760 and 1,140 ppm) inhibited the growth and development of *A. ostoyae* rhizomorphs initiated from inocula placed in quartz sand.
- The projected increase in CO<sub>2</sub> concentration in the atmosphere should not cause the future increase in the amount of *A. ostoyae* infectious material.

## Conflict of interest

The authors declare the lack of potential conflicts.

## Acknowledgements and source of funding

The research was financed from the funds of the Ministry of Science and Higher Education as part of a research project no. N309 019 31/2393.

The authors would like to express their gratitude to the manuscript reviewers, whose critical remarks helped to significantly improve the manuscript, as well as to Dr Joanna Ukalska for helping in performing statistical calculations and Mr. Kari Korhonen for providing testers to identify species of honey fungi.

## References

- Charkaborty S., von Tiedemann A., Teng P.S. 2000. Climate change: potential impact on plant diseases. *Environmental Pollution* 108: 317–326. DOI 10.1016/S0269-7491(99)00210-9.
- Climate Change 2001. Synthesis Report. Summary for Policymakers. An Assessment of the Intergovernmental Panel on Climate Change. UNFCCC, 2001.
- Fleischmann F., Raidl W.F., Osswald W.F. 2010. Changes in susceptibility of beech (*Fagus sylvatica*) seedlings towards *Phytophthora citricola* under the influence of elevated atmospheric CO<sub>2</sub> and nitrogen fertilization. *Environmental pollution* 158: 1051–1060. DOI 10.1016/j.envpol.2009.10.004.
- Garraway M.O. 1970. Rhizomorphs initiation and growth in *Armillaria mellea* promoted by o-aminobenzoic and p-aminobenzoic acids. *Phytopathology* 60: 861–865. DOI 10.1094/Phyto-60-861.
- Garraway M.O., Weinhold A.R. 1967. Ethanol-induced accumulation of a pentose in *Armillaria mellea*. *Canadian Journal of Microbiology* 12: 1705–1707. DOI 10.1139/m67-221.
- Garraway M.O., Weinhold A.R. 1968 a. Period of access to ethanol in relation to carbon utilization and rhizomorph initiation and growth in *Armillaria mellea*. *Phytopathology* 58: 1190–1191.
- Garraway M.O., Weinhold A.R. 1968 b. Influence of ethanol on the distribution of glucose C<sup>14</sup> assimilated by *Armillaria mellea*. *Phytopathology* 58: 1652–1657.
- Garrett K.A., Dendy S.P., Frank E.E., Rouse M.N., Travers S.E. 2006. Climate change effects on plant disease: genomes to ecosystems. *Annual Review of Phytopathology* 44: 489–509. DOI 10.1146/annurev.phyto.44.070505.143420.
- Guillaumin J.-J., Botton B. 2005. Ontogénie des armillaires. L'armillaire et le pourridié-agaric des végétaux ligneux. INRA, Paris, 151–175. ISBN 2-7380-1208-6.
- Guillaumin J.-J., Legrand P. 2005. Cycle infectieux de l'armillaire – Stratégie biologique des espèces européennes. L'armillaire et le pourridié-agaric des végétaux ligneux. INRA, Paris, 177–201. ISBN 2-7380-1208-6.
- Hartig R. 1873. Vorläufige Mitteilung über den Parasitismus von *Agaricus melleus* und dessen Rhizomorphen. *Botanische Zeitung* 31: 295–297.
- Harju A.M., Venalainen M., Anttonen S., Viitanen H., Kainulainen F., Vapaavuori E. 2003. Chemical factors affecting the brown-rot decay resistance of Scots pine hardwood. *Trees: Structure and Function* 17(3): 263–268. DOI 10.1007/s00468-002-0233-z.
- Hermes D.A., Mattson W.J. 1992. The Dilemma of Plants: to Grow or Defend. *Quarterly Review of Biology* 67(3): 283–335.
- Hintikka V. 1974. Notes on the ecology of *Armillaria mellea* in Finland. *Karstenia* 14: 12–31.
- Hintikka V., Korhonen K. 1970. Effects of carbon dioxide on the growth of lignicolous and soil-inhabiting Hymenomycetes. *Communicationes Instituti Forestalis Fenniae* 62: 1–22.
- Jacques-Felix M. 1967. Recherches morphologiques, anatomiques, morphogénétiques et physiologiques sur des rhizomorphes de champignons supérieurs et sur le déterminisme de leur formation. *Bulletin trimestriel de la. Societe mycologique de France* 83: 9–103.
- Jacques-Felix M. 1968. Recherches sur la morphogénese des rhizomorphes et télépodes en culture pure. *Bulletin trimestriel de la Societe mycologique de France* 84: 162–307.
- Korhonen K. 1978. Interfertility and clonal size in the *Armillaria mellea* complex. *Karstenia* 18: 31–42.
- Lech P., Żółciak A. 2006. Wzrost sadzonek sosny zwyczajnej i rozwój ryzomorf opieńki ciemnej w warunkach podwyższonej koncentracji CO<sub>2</sub> w powietrzu. *Leśne Prace Badawcze* 4: 17–34.
- Lech P., Żółciak A. 2017. Podatność sadzonek sosny zwyczajnej i świerka pospolitego na infekcje przez opieńkę ciemną w warunkach podwyższonych koncentracji CO<sub>2</sub> w powietrzu. *Sylvan* 161(5): 385–394.
- Lung-Escarmant B., Legrand P., Guillaumin J.-J., Guyon D. 2005. Épidémiologie.
- L'armillaire et le pourridié-agaric des végétaux ligneux. INRA, Paris, 203–219. ISBN 2-7380-1208-6.
- Łakomy P. 2004. Środowiskowe uwarunkowania zasiedlenia pniaków drzew liściastych przez wybrane gatunki grzybów saprotroficznych oraz grzyby rodzaju *Armillaria*. *Roczniki Akademii Rolniczej w Poznaniu. Rozprawy naukowe* 355: 1–164.

- Manning W.J., von Tiedemann A. 1995. Climate change: potential effects of increased atmospheric carbon dioxide (CO<sub>2</sub>), ozone (O<sub>3</sub>) and ultraviolet-B (UV-B) radiation on plant diseases. *Environmental pollution* 88: 219–245. DOI 10.1016/0269-7491(95)91446-R.
- Mańka K. 1953. Badania terenowe i laboratoryjne nad opieńką miodową - *Armillaria mellea* (Vahl.) Quél. *Prace Instytutu Badawczego Leśnictwa* 94: 1–96.
- Morrison D.J. 1975. Ion uptake by rhizomorphs of *Armillaria mellea*. *Canadian Journal of Botany* 53: 48–51. DOI 10.1139/b75-006.
- Morrison D.J. 1976. Vertical distribution of *Armillaria mellea* rhizomorphs in soil. *Transactions of the British Mycological Society* 66: 393–399. DOI 10.1016/S0007-1536(76)80207-0.
- Olszyk D.M., Johnson M.G., Phillips R.J., Seidler D.T., Tingey D.T., Watrud L.S. 2001. Interactive effects of CO<sub>2</sub> and O<sub>3</sub> on ponderosa pine plant/litter/soil mesocosm. *Environmental Pollution* 115(3): 447–462. DOI 10.1016/S0269-7491(01)00234-2.
- Oszako T., Sikora K., Borys M., Kubiak K., Tkaczyk M. 2016. *Phytophthora quercina* infections in elevated CO<sub>2</sub> concentrations. *Folia Forestalia Polonica, series A*, 58(3): 131–141. DOI 10.1515/ffp-2016-0015.
- Pentland G.D. 1965. Stimulation of rhizomorph development of *Armillaria mellea* by *Aureobasidium pullulans* in artificial culture. *Canadian Journal of Microbiology* 11: 345–350. DOI 10.1139/m65-042.
- Pentland G.D. 1967. Ethanol produced by *Aureobasidium pullulans* and its effect on the growth of *Armillaria mellea*. *Canadian Journal of Microbiology* 13: 1631–1639. DOI 10.1139/m67-213.
- Percy K.E., Awmack C.S., Lindroth R.I., Kubiske M.E., Kopper B.J., Isebrands J.G., Pregitzer K.S., Hendrey G.R., Dickson R.E., Zak D.R., Oksanen E., Sober J., Harrington R., Karnosky D.F. 2002. Altered performance of forest pests under atmospheres enriched by CO<sub>2</sub> and O<sub>3</sub>. *Nature* 420: 403–407. DOI 10.1038/nature01028.
- Przybył K. 1998. Effects of soil inorganic nutrients on rhizomorph growth by *Armillaria* sp. 9<sup>th</sup> International Conference of Root and Butt Rots of Forest Trees, str. 442. Carcans-Maubuisson (France), September 1-7, 1997. Ed. INRA, Paris, Abstr. ISBN 2-7380-0821-6.
- Redfern D.B. 1973. Growth and behaviour of *Armillaria mellea* rhizomorphs in soil. *Transactions of the British Mycological Society* 61: 569–581. DOI 10.1016/S0007-1536(73)80126-3.
- Redfern D.B., Filip G.M. 1991. Inoculum and infection, w: Shaw C.G., Kile G.A. (red.) *Armillaria Root Disease*. Agriculture Handbook, no. 691, U.S.D.A. Forest Service, 48–60.
- Rykowski K. 1984. Niektóre troficzne uwarunkowania patogeniczności *Armillaria mellea* (Vahl) Quél. w uprawach sosnowych. *Prace Instytutu Badawczego Leśnictwa, seria A* 640: 1–140.
- Schinner F., Concin R. 1981. Carbon dioxide fixation by wood rotting fungi. *European Journal of Forest Pathology* 11(1–2): 120–123. DOI 10.1111/j.1439-0329.1981.tb00077.
- Sortkjaer O., Allermann K. 1972 a. Rhizomorph formation in fungi. I. Stimulation by ethanol and acetate and inhibition by Disulfiram of growth and rhizomorph formation in *Armillaria mellea*. *Physiologia Plantarum* 26: 376–380. DOI 10.1111/j.1399-3054.1972.tb01125.x.
- Sortkjaer O., Allermann K. 1972 b. Rhizomorph formation in fungi. II. The effect of 12 different alcohols on growth and rhizomorph formation in *Armillaria mellea* and *Clitocybe geotropa*. *Physiologia Plantarum* 28: 51–55. DOI 10.1111/j.1399-3054.1972.tb01125.x.
- Sortkjaer O., Allermann K. 1972 c. Rhizomorph formation in fungi. III. The effect of ethanol on the synthesis of DNA and RNA and uptake of Asparagine and phosphate in *Armillaria mellea*. *Physiologia Plantarum* 29: 129–133. DOI 10.1111/j.1399-3054.1972.tb01125.x.
- Tkaczyk M., Sikora K., Nowakowska J.A., Kubiak K., Oszako T. 2014. Effects of CO<sub>2</sub> enhancement on beech (*Fagus sylvatica* L.) seedling root rot due to *Phytophthora plurivora* and *Phytophthora cactorum*. *Folia Forestalia Polonica, series A* 56(3): 149–156. DOI 10.2478/ffp-2014-0016.
- Żółciak A. 1999. Identyfikacja gatunków grzybów z rodzaju *Armillaria* (Fr.: Fr.) Staude w Polsce. *Prace Instytutu Badawczego Leśnictwa, seria A* 888: 3–19.
- Żółciak A. 1999. Występowanie grzybów z rodzaju *Armillaria* (Fr.: Fr.) Staude w kompleksach leśnych w Polsce. *Prace Instytutu Badawczego Leśnictwa, seria A* 890: 29–40.
- Żółciak A. 2005. Opieńki. CILP, Warszawa, 1–111. ISBN 8388478-85-0.

## Authors' contribution

P.L. – research concept, literature review, methodology, measurements, preparation of test results, statistical analysis, manuscript writing/editing; A.Ż. – collection/identification of fungal material in the field, preparation of inoculum, methodology, literature review, manuscript writing/editing.