

Potentially invasive fungal pathogens in forests of Nepal

Sanjay Kumar Jha¹, Hari Sharan Adhikari² ✉, Lal Bahadur Thapa¹, Sunita Ulak³,
Dhirendra Kumar Pradhan⁴, Yam Prasad Pokhare⁵, Shambhu Kumar⁵

¹ Tribhuvan University, Institute of Science and Technology, Central Department of Botany, Kathmandu, Nepal

² Tribhuvan University, Institute of Science and Technology, Amrit Campus, Department of Botany, Kathmandu, Nepal,
e-mail: aharisharan@gmail.com

³ Ministry of Forest and Environment, Forest Research and Training Center (FRTC), Babarmahal, Kathmandu, Nepal

⁴ Ministry of Forest and Environment, Forest and Watershed Division, Singh Durbar, Kathmandu, Nepal

⁵ Kerala Forest Research Institute, Peechi, Kerala, India

ABSTRACT

Potentially invasive fungal pathogens of tree species in Nepal were isolated and identified in this study. A survey was conducted throughout the country, and the infected parts of different host trees were cultured in potato dextrose agar medium to isolate the pathogenic fungi. Altogether, 10 fungal pathogens were identified as causing foliar and trunk diseases. Two of the pathogens, *Nectria* sp. (stem canker) and *Phytophthora alni* subsp. *alni* (bleeding canker), were reported from the stems and trunks of *Alnus nepalensis* and *Shorea robusta*, respectively. The pathogens *Neonectria neomacrospora* (needle cast), *Aureobasidium apocryptum* (anthracnose), *Golovinomyces cichoracearum* (powdery mildew), *Calonectria reteaudii* (blight), *Dothistroma septosporum* (needle cast), *Calonectria indusiata* (spot), *Olivea tectonae* (rust) and *Rhytisma acerinum* (tar spot) were isolated from the foliar part of the host trees *Abies spectabilis*, *Acer laevigatum*, *Alnus nepalensis*, *Eucalyptus camaldulensis*, *Pinus roxburghii*, *Rhododendron arboreum*, *Tectona grandis* and *Toona ciliata*, respectively. These species could be the potential invasive forest pathogens in Nepal. An effective long-term future plan needs to be developed for managing these forest diseases to maintain the forest health of the country.

KEY WORDS

forest disease, forest health, fungal pathogens, invasive pathogens, plant-microbe interaction

INTRODUCTION

Forests, vital ecosystems, provide a wide range of ecosystem services, such as timber and non-timber forest products, carbon sequestration, biodiversity conservation, soil formation and nutrient cycling. All these services can be achieved from healthy forests. A healthy forest refers to the overall condition of the forest and its ability

to support ecological and economic functions over time (Trumbore et al. 2015). Healthy forests are not only free of diseases, pest infestations, alien invasions and pollution, but also resistant to a wide variety of stresses and diseases (Warren 2007; Linnakoski et al. 2019).

Currently, climate change events, pollution and invasions of alien species have increased the risk of diseases in forests, as these factors weaken plants, making

them more susceptible to diseases and pests (Solomou et al. 2019; Burdon and Zhan 2020; Panzavolta et al. 2021). Various types of diseases caused by fungi, bacteria, viruses, nematodes, insects and other pests have become more severe with increasing intensities of the abovementioned factors; therefore, forest health is deteriorating and productivity has been reduced (Guégan et al. 2020; El-Sayed and Kamel 2020; Paseka et al. 2020).

Nepal is a country with diverse forest ecosystems from tropical lowlands to subalpine regions that provide a range of ecosystem services and support livelihoods for many communities (Lamsal et al. 2018). As climate change events, pollution and alien plant invasion have been serious problems, Nepal's forests are facing a range of direct negative impacts from these factors (Thapa et al. 2017, 2020; Shrestha and Shrestha 2021; Charmakar et al. 2022; Wani et al. 2023). In addition, serious infestation by pathogens has been reported throughout the country (Malla and Pokharel 2018). For-

est disease is an important and complex area of research that requires interdisciplinary approaches to understand the causes and consequences of diseases in forests, but it is often a neglected area of research in Nepal. This study was carried out with the aim of assessing highly problematic diseases of tree species in Nepal. The study will have significance in understanding and will be useful for future planning of the management of forest pathogens and preserving forest health in the country.

MATERIAL AND METHODS

Collection of disease sample of plants

In this study, a total of seven districts (Taplejung, Nawalpur, Tanahun, Lamjung, Dang, Dadheldhura and Darchula) of eastern, central and western Nepal were visited (Fig. 1, Tab. 1). Diseased plant samples were collected during August–September 2020, and the samples were

packaged in sterile paper bags with labelling of latitude, longitude, altitude and collection number. The samples were kept in icebox and transported to the Plant Pathology Laboratory of Forest Research and Training Center, Ministry of Forestry, Babarmahal, Kathmandu, Nepal. The samples were stored in refrigerator at 4°C for further analysis. All the host specimens were also identified using standard literature such as Shrestha et al. (2022).

Isolation and culturing of phytopathogens

The diseased samples were washed thoroughly, surface sterilised with 0.2% sodium hypochlorite and again washed with sterile distilled water. Then, small fragments (5 × 5 mm) of infected portions were placed on Petri plates containing potato dextrose agar media supplemented with 100 mg/L streptomycin sulphate. The plates were incubated at 25±2°C for 5–7 days (Johnston and Booth 1983). The hyphal tips from each developing colony's margin were subcultured as pure

Table 1. Details of sample collection sites in Nepal

S. No.	Name of host plant	Diseased parts	Location	Elevation (m)	Forest	Municipality and district
1	<i>Abies spectabilis</i>	needles	27.422°N 87.763°E	3413	Mayam Patal CF	Phungling Municipality, Taplejung
2	<i>Acer laevigatum</i>	leaves	27.365°N 87.727°E	2642	Deurali-Bhitri CF	
3	<i>Rhododendron arboretum</i>	leaves	27.427°N 87.766°E	364	Mayam Patal CF	
4	<i>Alnus nepalensis</i>	leaves	28.293°N 84.369°E	901	Tatopani CF	Marsyangdi RM, Lamjung
5	<i>Alnus nepalensis</i>	trunk	29.336°N 80.606°E	1639	Ugratara CF	Ajaymeru RM, Dadheldhura
6	<i>Eucalyptus camaldulensis</i>	leaves	28.095°N 80.199°E	561	Eucalyptus PF	Dangisharan RM, Dang
7	<i>Pinus roxburghii</i>	needles	29.698°N 80.544°E	1416	Shalya Shikhar CF	Shailyashikhar Municipality, Darchula
8	<i>Shorea robusta</i>	stem	27.622°N 84.056°E	197	Sal SF	Madhyabindu Municipality, Nawalpur
9	<i>Tectona grandis</i>	leaves	28.095°N 82.199°E	561	Teak PF	Dangisharan RM, Dang
10	<i>Toona ciliata</i>	leaves	27.989°N 84.268°E	487	Gairigau CF	Vyas Municipality, Tanahun

Note. CF: Community Forest, PF: Planted Forest, SF: Scientific Forest, RM: Rural Municipality



Figure 1. Map showing sample collection sites

isolates. Non-culturable pathogens were observed by cutting the leaf sections through the infected portions.

Identification of phytopathogens

Cultured pathogens were observed using an OPTIKA B-380 LED light microscope. In addition, measurements of isolated fungi were taken using OPTIKA Microscopy software. The pathogens were identified based on the morphological characteristics of colonies, hyphae, fruiting bodies and reproductive structures, following Gilman (1957), Barnett and Hunter (1998), Braun and Cook (2012) and Campbell and Johnson (2013). Taxonomic verification and assigning the nomenclature of pathogens and their current names were performed based on Mycobank (<https://www.mycobank.org/>).

RESULTS AND DISCUSSION

Altogether, 10 pathogenic fungi from different tree hosts were isolated and identified. Among them, eight pathogens were identified from foliar diseases and two were identified from stem and trunk diseases. The foliar pathogens were *Neonectria neomacrospora*,

pora from the host *Abies spectabilis*, *Aureobasidium apocryptum* from *Acer laevigatum* and *Golovinomyces cichoracearum* from *Alnus nepalensis*. Similarly, *Calonectria reteaudii* was isolated from the leaves of *Eucalyptus camaldulensis*, *Dothistroma septosporum* from *Pinus roxburghii* needles and *Calonectria indusiata* from *Rhododendron arboreum*. The pathogens *Olivea tectonae* and *Rhytisma acerinum* were isolated from the host trees *Tectona grandis* and *Toona ciliata*, respectively. Meanwhile, *Phytophthora alni* subsp. *Alni* and *Nectria* sp. were isolated from the stems of *Alnus nepalensis* and *Shorea robusta*, respectively (Annex I). The diseases caused by these pathogens, symptomatology and identifying characteristics are given in Table 2.

The distribution of fungal pathogens is shown in Figure 2. The pathogens *Neonectria neomacrospora*, *Aureobasidium apocryptum* and *Calonectria indusiata* were reported from the eastern part of Nepal (Koshi Province). From the western part (Gandaki Province), the pathogens *Golovinomyces cichoracearum*, *Nectria* sp. and *Rhytisma acerinum* were reported. Similarly, two of the pathogens (*Calonectria reteaudii* and *Olivea tectonae*) were found in western Nepal (Lumbini Province) and *Dothistroma septosporum* and *Phytophthora*

Table 2. Host plants, disease name, pathogens, symptomatology and identifying characters

S. No.	Host plant	Disease	Pathogen identified	Symptomatology	Identifying characters	
					colony colour	vegetative and reproductive characters
1.	<i>Abies spectabilis</i>	needle cast	<i>Neonectria neomacrospora</i>	infected needles were light brown to dark brown; young needles were uninfected	white	<i>hyphae</i> : septate; <i>microconidia</i> : size 10.3–13.6 × 3.7 µm, multicellular, cylindrical or elongated, slightly curved and up to 93.11 µm
2.	<i>Acer laevigatum</i>	leaf anthracnose	<i>Aureobasidium apocryptum</i>	light brown spots and large necrotic patches on leaves, some of the leaves were entirely covered by necrotic patches	greyish white	<i>hyphae</i> : septate; small acervuli erupted through the epidermis of leaf; <i>conidia</i> : size 6.5–10.5 × 3–4.5 µm, hyaline, non-septate borne on short, broad conidiophores
3.	<i>Alnus nepalensis</i>	powdery mildew	<i>Golovinomyces cichoracearum</i>	white powdery spores on leaves, chlorosis and premature dropping	white	<i>hyphae</i> : hyaline and smooth (3.5–6.8 µm wide); <i>conidia</i> : hyaline, cylindrical or ellipsoidal, size 28.7–18.7 µm, developed at the apex of conidiophores
4.	<i>Alnus nepalensis</i>	bleeding canker	<i>Phytophthora alni</i> subsp. <i>alni</i>	the infected trunk was oozing reddish brown fluid from cracks, dieback symptom was also reported	light brown, white at maturity	<i>hyphae</i> : aseptate; <i>sporangiophores</i> : slender, unbranched; <i>zoosporangia</i> : size 35–65 × 24–50 µm, ovoid or pear shape
5.	<i>Eucalyptus camaldulensis</i>	leaf blight	<i>Calonectria reteaudii</i>	grey water-soaked spots, necrotic patches, some leaves were defoliated	light grey	<i>hyphae</i> : septate; <i>conidia</i> : size 65–85 × 5–6 µm, straight or cylindrical, round at both ends
6.	<i>Pinus roxburghii</i>	pinus needle cast	<i>Dothistroma septosporum</i>	yellowish-tan spots on needles, reddish brown bands around the needle, premature death and drop of infected needles	white with abundant aerial hyphae	<i>Conidiophores</i> : numerous, hyaline; <i>conidia</i> : septate, hyaline, straight or slightly curved, size 16.4–23.6 × 3 µm
7.	<i>Rhododendron arboreum</i>	leaf spot	<i>Calonectria indusiata</i>	dark brown spots on leaves, somewhere the spots covered whole leaves, premature defoliation	white with abundant aerial hyphae	<i>hyphae</i> : septate; <i>conidiophores</i> : penicillate, two to six phialides; <i>conidia</i> : size 60–70 × 4–6 µm, straight or cylindrical, round at both ends, septate
8.	<i>Shorea robusta</i>	stem canker	<i>Nectria</i> sp.	oval to elongated, brown to reddish lesions on the bark of trunk and branches, oozing sap with foul smell	cottony with white with aerial hyphae	<i>Hyphae</i> : septate; <i>conidiophores</i> : verticillate, one to three branched; <i>conidia</i> : ellipsoidal to cylindrical or slightly curved, hyaline, smooth, rounded at both ends
9.	<i>Tectona grandis</i>	leaf rust	<i>Olivea tectonae</i>	small, angular, brown to grey necrotic areas on upper leaf surface, necrosis due to coalesced lesions, powdery yellowish rusts	not cultured	<i>Uredospores</i> : powdery, orange coloured, ovoid and echinulated, size 18–23 × 17.5–24.8 µm
10.	<i>Toona ciliata</i>	tar spot	<i>Rhytisma acerinum</i>	small, superficial black tar-like spots	greyish brown	<i>ascmata</i> : 40–120 µm wide with asci and paraphyses; <i>ascospores</i> : ellipsoidal, blunt at both ends, size 40–70 × 9–10 µm

alni subsp. *alni* were reported from the far western part of Nepal (Sudurpashchim Province) (Fig. 2).

The fungal pathogen *Neonectria neomacrospora* was reported in Norway, Denmark and England from 2008 to




Figure 2. Distribution of potential invasive fungal pathogens in Nepal

2015, causing stem canker and dieback disease in different species of *Abies* (Nielsen et al. 2017). Due to the re-emergence of this pathogen causing different symptoms on shoots, stems, branches, needles and cones, it was identified as an aggressive pathogen of *Abies* and was included in the European and Mediterranean Plant Protection Organization (EPPO) Alert list (González et al. 2021). In Nepal, the pathogen causing needle cast disease was reported in *Abies spectabilis* by González et al. (2021).

Another pathogen, *Aureobasidium apocryptum*, was reported in the leaves of *Acer* sp. from Canada (Creelman 1965) and North America (Vujanovic and Brisson 2002). Bamadhaj et al. (2015) reported this pathogen as a potential invasive pathogen. The incidence of this pathogen in Nepal might be a new record from Nepal. Literature related to the pathogen *Golovinomyces cichoracearum* in *Alnus nepalensis* was not found. Instead, *Erysiphe penicillata* and *Oidium* sp. were reported in *Alnus glutinosa* (Mieslerová et al. 2020) and *Alnus nepalensis* (Srivastava and Verma 1990) from the Czech Republic and Sikkim (India), respectively. Another species of *Golovinomyces* was reported as an invasive pathogen by Félix-Gastelum et al. (2019) in Mexico.

A destructive pathogen *Phytophthora* from *Alnus* trees was found in 1993 in the United Kingdom and

was later confirmed to exist in various regions of Europe (Brasier et al. 1995). The pathogen was found to be locally very damaging and spreading with a threat to both natural and managed alder populations in Europe, Asia and North America (Redondo et al. 2015; Jung and Blaschke 2004; Sims et al. 2015). In Vietnam, *Phytophthora* sp. close to *Phytophthora tropicalis* was reported by Jung et al. (2020) from *Alnus nepalensis* causing bleeding canker and was designated *Phytophthora* sp. *tropicalis*-like 2. The pathogens isolated in this study were identified as *Phytophthora alni* subsp. *alni* on the basis of the morphological characteristics of the pathogen as described by Hansen (2012).

Calonectria reteaudii (Anamorph: *Cylindrocladium reteaudii*) is a well-recognised pathogen responsible for the decline of *Eucalyptus* trees (Bose et al. 2022). In Vietnam and Indonesia, 10 undescribed new species of *Calonectria* were reported in *Eucalyptus* and *Acacia* (Pham et al. 2019). Members of this complex are widespread worldwide in natural and *Eucalyptus* plantations and have recently been reported from India, Malaysia, southern China, Brazil and Thailand (Jessadarom et al. 2018; Wu and Chen 2021; Pham et al. 2022; Bose et al. 2022; Sanchez-Gonzalez et al. 2022). No previous literature was found regarding this pathogen in Nepal, and

therefore, it is expected that this is a new report from the country.

Dothistroma needle blight (DNB) is one of the most serious foliar diseases of *Pinus* spp. in many countries (Drenkhan et al. 2016). *Dothistroma septosporum* is the root cause of DNB, together with *Dothistroma pini* (Barnes et al. 2004). The disease symptoms for both pathogens are necrotic lesions on the needle, frequently accompanied by red bands, and early needle loss (Barnes et al. 2004). The disease caused by *Dothistroma septosporum* (reported from this study) in *Pinus* spp. has gained worldwide attention because of the terrible damage to pine plantations (Bakshi and Singh 1968; Barnes et al. 2022). Another species, *Calonectria*, is frequently associated with a variety of disease symptoms, such as leaf spot, stem rot, canker, blight, root and pod rot (Crous 2006). *Calonectria indusiata* was first reported in *Rhododendron* from Florida, USA (Crous et al. 2000) and described by Crous (2002) as the most prevalent canker disease pathogen.

Nectria canker is common in hardwood tree species from the temperate regions of the northern hemisphere (Lortie 1974; Yang et al. 2018). In the present research, canker in *Shorea robusta* was found to be caused by *Nectria* sp., but no literature was found supporting this finding. The pathogen *Olivea tectonae* is the major rust fungus in the tree species *Tectona grandis*, causing premature defoliation and reduction of the expected growth rates from nursery to adult plants (Cabral et al. 2010; Pathak et al. 2015; Koffi et al. 2018). This fungus is categorised as an invasive pathogen in the Oceania region, and the disease has also been reported throughout Asia (CABI/EPPO 2015; EPPO 2022).

Tar spot is another common plant disease caused by different pathogenic fungi depending on the type of host plant. Among the different causal agents, *Rhytisma acerinum* is the major agent that chiefly causes the disease in Mapple (Hudler et al. 1987), but it was also reported in *Toona ciliata* in the present research work, which was first reported in India (Chandel and Kumar 2017). The majority of pathogens were previously documented from different regions of the world (Farr et al. 2021); however, there was no prior record of these pathogens in Nepal.

Nepal's forests serve as an ecological bridge between South Asian and East Asian biomes, so pathogens discovered here pose immediate cross-border threats to neighbouring India, China and Bhutan. As

a sentinel site, Nepal's initial reports of these globally devastating infections serve as an early warning system for future continent-scale shifts, illustrating how invasive fungi might exploit new hosts and habitats. Climate change threatens to broaden disease ranges; rising temperatures may allow subtropical dangers to spread to temperate zones around the world, transforming localised infections into international crises (Gullino et al. 2022; Kumar and Mukhopadhyay 2025). Biocontrol presents an effective alternative to chemical pesticides and fertilisers in fighting forest pathogens, though more research is needed to advance biological control agents against such invasive threats (Balla et al. 2021).

CONCLUSIONS

Overall, a total of 10 pathogenic fungi that are potentially invasive pathogens from different host trees in Nepal were isolated and identified. Most of the pathogens caused foliar disease with different symptomatology, including blight, spots, powdery mildew, etc. The pathogens were prevalent in community-managed and planted forests. The prevalence of these pathogens was mostly on timber trees, such as *Shorea robusta*, *Abies spectabilis* and *Tectona grandis*. These findings constitute the first documented occurrences of these phytopathogens within Nepal, indicating the emergence of invasive forest diseases in the region. Critically, the detection of globally destructive agents such as *Phytophthora alni* subsp. *alni* (on *Shorea robusta*) and *Dothistroma septosporum* (on *Pinus roxburghii*), underscores their potential as transboundary phytosanitary threats to ecologically pivotal tree species. Consequently, comprehensive longitudinal surveillance and expanded research across Nepal's forest ecosystems are imperative to map pathogen distribution, assess invasion pathways and implement pre-emptive management strategies to mitigate further dissemination.

ACKNOWLEDGEMENTS

This work was supported by the FAO Forest Health project no. TCP/NEP/3702. The authors are also thankful to the Kerala Forest Research Institute (KFRI), Kerala, India, for technical support. Dr. K.V. Sankaran (TCDC

Expert, FAO), Dr. Binod Saha (FAO Nepal), Dr. Shiro Sathiyapala (FAO), Dr. Ilias Animon (FAO- RAP) and Prof. Dr. Bharat Babu Shrestha (Central Department of Botany, Tribhuvan University, Nepal) are acknowledged for their valuable suggestions and all kinds of support.

REFERENCES

- Agrios, G.N. 2005. Plant Pathology. Elsevier, Burlington, MA, USA, 473–483.
- Bakshi, B.K., Singh, S. 1968. Dothistroma blight – a potential threat to *Pinus radiata* plantations in India. *Indian Forester*, 94 (11), 824–825.
- Balla, A. et al. 2021. The threat of pests and pathogens and the potential for biological control in forest ecosystems. *Forests*, 12 (11), 1579.
- Bamadhaj, H.M., Chan, G.F., Rashid, N.A.A. 2015. Aureobasidium. In: Molecular biology of food and water borne mycotoxigenic and mycotic fungi (eds. R. Russell M. Paterson, Nelson Lima). CRC Press, Boca Raton. DOI: <https://doi.org/10.1201/b18645>.
- Barnes, I., Crous, P.W., Wingfield, B.D., Wingfield, M.J. 2004. Multigene phylogenies reveal that red band needle blight of *Pinus* is caused by two distinct species of *Dothistroma*, *D. septosporum* and *D. pini*. *Studies in Mycology*, 50 (2), 551–565.
- Barnes, I., van der Nest, A., Granados, G.M., Wingfield, M.J. 2022. Dothistroma needle blight. In: Forest microbiology (eds. F. Asiegbu, A. Kovalchuk). Academic Press, London, 179–199.
- Barnett, H.L., Hunter, B. 1998. Illustrated genera of imperfect fungi. APS Press, Minnesota, USA.
- Bose, R., Banerjee, S., Negi, N., Pandey, A., Bhandari, M.S., Pandey, S. 2022. Identification and pathogenicity of *Calonectria pseudoreteauidii* causing leaf blight of Eucalyptus – a new record for India. *Physiological and Molecular Plant Pathology*, 122, 101917.
- Brasier, C.M., Rose, J., Gibbs, J.N. 1995. An unusual *Phytophthora* associated with widespread alder mortality in Britain. *Plant Pathology*, 44 (6), 999–1007.
- Braun, U. 2012. Taxonomic manual of Erysiphales (powdery mildews). CBS Biodiversity series 11. CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands.
- Burdon, J.J., Zhan, J. 2020. Climate change and disease in plant communities. *PLoS Biology*, 18 (11), e3000949.
- CABI/EPPO. 2015. *Olivea tectonae*. [Distribution map]. In: Distribution maps of plant diseases. CABI, Wallingford, UK.
- Cabral, P.G.C. et al. 2010. First report of teak leaf rust disease caused by *Olivea tectonae* in Brazil. *Australasian Plant Disease Notes*, 5 (1), 113–114.
- Campbell, C.K., Johnson, E.M. 2013. Identification of pathogenic fungi. John Wiley & Sons, Chichester.
- Chandel, S., Kumar, V. 2017. First report of tar spot of Toona (*Toona ciliata*) in India. *Journal of Applied and Natural Science*, 9 (2), 784–785.
- Charmakar, S. et al. 2022. Invasive alien plants, insect pests and pathogens in planted and natural forests in Nepal: Key lessons from an online survey on distribution and impacts. In: Proceeding of XV World Forestry Congress, 2–6 May 2022, Seoul, South Korea, 1–8.
- Creelman, D.W. 1964. Summary of the prevalence of plant diseases in Canada in 1963. *Canadian Plant Disease Survey*, 44 (1), 1–82.
- Crous, P.W. 2002. Taxonomy and pathology of *Cylindrocladium* (*Calonectria*) and allied genera. American Phytopathological Society (APS Press), St. Paul, MN, USA.
- Crous, P.W., Groenewald, J.Z., Risede, J.M., Simoneau, P., Hyde, K.D. 2006. *Calonectria* species and their *Cylindrocladium* anamorphs: Species with clavate vesicles. *Study of Mycology*, 55, 213–226.
- Crous, P.W., Kang, J.C., Schoch, C.L., Mchau, G.R. 2000. Phylogenetic relationships of *Cylindrocladium pseudogratile* and *Cylindrocladium rumohrae* with morphologically similar taxa, based on morphology and DNA sequences of internal transcribed spacers and beta-tubulin. *Canadian Journal of Botany*, 77 (12), 1813–1820.
- Drenkhan, R. et al. 2016. Global geographic distribution and host range of *Dothistroma* species: a comprehensive review. *Forest Pathology*, 46 (5), 408–442.
- El-Sayed, A., Kamel, M. 2020. Climatic changes and their role in emergence and re-emergence of diseases. *Environmental Science and Pollution Research*, 27, 22336–22352.

- EPPO. 2022. EPPO Global Database. Paris, France. Available at <https://gd.eppo.int/>.
- Farr, D.F., Rossman, A.Y., Castlebury, L.A. 2021. United States National Fungus Collections Fungus-Host Dataset. Ag Data Commons. DOI: <https://doi.org/10.15482/USDA.ADC/1524414>.
- Félix-Gastélum, R. et al. 2019. Powdery mildew caused by *Golovinomyces spadicus* on wild sunflower in Sinaloa, Mexico. *Canadian Journal of Plant Pathology*, 41 (2), 301–309.
- Freer-Smith, P.H., Webber, J.F. 2017. Tree pests and diseases: the threat to biodiversity and the delivery of ecosystem services. *Biodiversity and Conservation*, 26 (13), 3167–3181.
- Geng, X., Liu, Y., Li, J., Li, Z., Shu, J., Wu, G. 2022. Identification and characterization of *Nectria pseudotrichia* associated with camellia canker disease in China. *Forests*, 13 (1), 29.
- Gilman, J. 1957. A manual of soil fungi. *Soil Science*, 84 (2), 183.
- Gonzalez, M. et al. 2021. Incidence of the emerging pathogen *Neonectria neomacrospora* on *Abies* taxa in the National Arboreta in England (UK). *Forest Ecology and Management*, 492, 119207.
- Guégan, J.F., Ayoub, A., Cappelle, J., De Thoisy, B. 2020. Forests and emerging infectious diseases: unleashing the beast within. *Environmental Research Letters*, 15 (8), 083007.
- Gullino, M. L. et al. 2022. Climate change and pathways used by pests as challenges to plant health in agriculture and forestry. *Sustainability*, 14 (19), 12421.
- Hansen, E.M. 2012. *Phytophthora alni*. *Forest Phytophthoras*, 2 (1).
- Hudler, G.W., Banik, M.T., Miller, S.G. 1987. Unusual epidemic of tar spot on Norway maple in upstate New York. *Plant Disease*, 71 (1), 65–67.
- Inghelbrecht, S., Gehesquière, B., Heungens, K. 2011. First report of *Calonectria* leaf spot caused by *Calonectria colhounii* (anamorph *Cylindrocladium colhounii*) on *Rhododendron* in Belgium. *Plant disease*, 95 (11), 1477–1477.
- IPPC. 2015. Detection of teak leaf rust. IPPC Official Pest Report, No. AUS-01/3. FAO, Rome, Italy.
- Jactel, H., Koricheva, J., Castagnyrol, B. 2019. Responses of forest insect pests to climate change: not so simple. *Current Opinion in Insect Science*, 35, 103–108.
- Jessadarom, H., Phetruang, W., Haitook, S., Cheewangkoon, R. 2018. Isolation of *Calonectria sulawesensis* from soil in Thailand and its pathogenicity against *Eucalyptus camaldulensis*. *Plant Pathology and Quarantine*, 8, 1–8.
- Johnston, A., Booth, C. (eds.) 1983. Plant Pathologist's Pocketbook. CAB, Slough, UK.
- Jung, T., Blaschke, M. 2004. Phytophthora root and collar rot of alders in Bavaria: distribution, modes of spread and possible management strategies. *Plant pathology*, 53(2), 197–208.
- Jung, T. et al. 2020. A survey in natural forest ecosystems of Vietnam reveals high diversity of both new and described *Phytophthora* taxa including *P. ramorum*. *Forests*, 11 (1), 93.
- Koffi, N.B.C., N'douba, A.P., Kouassi, C.K., Koffi, A.M.H., Diallo, H.A. 2018. First occurrence of teak leaf rust caused by *Olivea tectonae* in Cote d'Ivoire and Africa. *New Disease Reports*, 22.
- Kumar, D., Mukhopadhyay, R. 2025. Climate change and plant pathogens: Understanding dynamics, risks and mitigation strategies. *Plant Pathology*, 74 (1), 59–68.
- Lamsal, P., Kumar, L., Atreya, K., Pant, K.P. 2018. Forest ecosystem services in Nepal: a retrospective synthesis, research gaps and implications in the context of climate change. *International Forestry Review*, 20 (4), 506–537.
- Linnakoski, R., Kasanen, R., Dounavi, A., Forbes, K.M. 2019. Forest health under climate change: effects on tree resilience, and pest and pathogen dynamics. *Frontiers in Plant Science*, 10, 1157.
- Lortie, M. 1974. *Nectria* canker of hardwoods. URL: <https://agris.fao.org/agris-search/search.do?recordID=US201300578159>.
- Malla, R., Pokharel, K.K. 2018. Forest pests and pathogens problem in different forest types of Nepal. Department of Forest Research and Survey, Kathmandu, Nepal.
- Mesquita, J.B., Santos, Í., Ribeiro, G.T., dos Santos, M.J.C. 2016. Rust (*Olivea neotectonae*) occurrence on teak plants in Sergipe, Brazil. *Summa Phytopathologica*, 42 (3), 278–279.
- Mieslerová, B., Sedlářová, M., Michutová, M., Petřeková, V., Cook, R., Lebeda, A. 2020. Powdery mildews on trees and shrubs in botanical gardens,

- parks and urban green areas in the Czech Republic. *Forests*, 11 (9), 967.
- Nielsen, U.B., Xu, J., Nielsen, K.N., Talgø, V., Hansen, O.K., Thomsen, I.M. 2017. Species variation in susceptibility to the fungus *Neonectria neomacropora* in the genus *Abies*. *Scandinavian Journal of Forest Research*, 32 (5), 421–431.
- Panzavolta, T., Bracalini, M., Benigno, A., Moricca, S. 2021. Alien invasive pathogens and pests harming trees, forests, and plantations: Pathways, global consequences and management. *Forests*, 12 (10), 1364.
- Paseka, R.E. et al. 2020. Disease-mediated ecosystem services: Pathogens, plants, and people. *Trends in Ecology and Evolution*, 35 (8), 731–743.
- Pathak, H., Maru, S., Satya, H.N., Silawat, S.C. 2015. Fungal diseases of trees in forest nurseries of Indore, India. *Journal of Plant Pathology and Microbiology*, 6 (8), 297.
- Pérez, M., López, M.O., Martí, O. 2009. *Olivea tectonae*, leaf rust of teak, occurs in Cuba. *Plant Pathology*, 58 (2), 397.
- Pham, N.Q. et al. 2019. Ten new species of *Calonectria* from Indonesia and Vietnam. *Mycologia*, 111 (1), 78–102.
- Pham, N.Q., Marincowitz, S., Chen, S., Yaparudin, Y., Wingfield, M.J. 2022. *Calonectria* species, including four novel taxa, associated with *Eucalyptus* in Malaysia. *Mycological Progress*, 21 (1), 181–197.
- Redondo, M.A., Boberg, J., Olsson, C.H., Oliva, J. 2015. Winter conditions correlate with *Phytophthora alni* subspecies distribution in Southern Sweden. *Phytopathology*, 105 (9), 1191–1197.
- Roy, B.A. et al. 2014. Increasing forest loss worldwide from invasive pests requires new trade regulations. *Frontiers of Ecology and Environment*, 12, 457–465.
- Sanchez-Gonzalez, E.I., Soares, T.D.P.F., Zarpelon, T.G., Zauza, E.A.V., Mafia, R.G., Ferreira, M.A. 2022. Two new species of *Calonectria* (Hypocreales, Nectriaceae) causing *Eucalyptus* leaf blight in Brazil. *MycKeys*, 91, 169–197.
- Shrestha, B.B., Shrestha, K.K. 2021. Invasions of alien plant species in Nepal: Patterns and process. In: Invasive alien species: observations and issues from around the world (eds. T. Pullaiah, M.R. Lelmin). John Wiley and Sons Ltd, 168–183.
- Shrestha, K.K., Bhandari, P., Bhattarai, S. 2022. Plants of Nepal (Gymnosperms and Angiosperms). Heritage Publishers and Distributors Pvt. Limited, Kathmandu, Nepal.
- Sims, L.L., Sutton, W., Reeser, P., Hansen, E.M. 2015. The *Phytophthora* species assemblage and diversity in riparian alder ecosystems of western Oregon, USA. *Mycologia*, 107 (5), 889–902.
- Solomou, A.D., Topalidou, E.T., Germani R., Argiri, A., Karetos, G. 2019. Importance, utilization and health of urban forests: A review. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 47 (1), 10–16.
- Srivastava, L.S., Verma, R.N. 1990. Some newly recorded fungal diseases of forest trees of Sikkim. *Indian Forester*, 116 (1), 89–90.
- Thapa, L.B., Kaewchumnong, K., Sinkkonen, A., Sridith, K.J.W.R. 2017. Plant invasiveness and target plant density: high densities of native *Schima wallichii* seedlings reduce negative effects of invasive *Ageratina adenophora*. *Weed Research*, 57 (2), 72–80.
- Thapa, L.B., Kaewchumnong, K., Sinkkonen, A., Sridith, K. 2020. “Soaked in rainwater” effect of *Ageratina adenophora* on seedling growth and development of native tree species in Nepal. *Flora*, 263, 151554.
- Trumbore, S., Brando, P., Hartmann, H. 2015. Forest health and global change. *Science*, 349 (6250), 814–818.
- Vujanovic, V., Brisson, J. 2002. A comparative study of endophytic mycobiota in leaves of *Acer saccharum* in eastern North America. *Mycological Progress*, 1 (2), 147–154.
- Wani, Z.A., Fatima, T., Verma, O., Sharma, V., Pant, S. 2023. Impacts of climate change on plants with special reference to the Himalayan region. In: Climate change adaptation, risk management and sustainable practices in the Himalaya (eds. S. Sharma, J.C. Kuniyal, P. Chand, P. Singh). Springer International Publishing, Cham, 237–251.
- Warren, W.A. 2007. What is a healthy forest? Definitions, rationales, and the lifeworld. *Society and Natural Resources*, 20 (2), 99–117.
- Wu, W., Chen, S. 2021. Species diversity, mating strategy and pathogenicity of *Calonectria* species from diseased leaves and soils in the *Eucalyptus* plantation in southern China. *Journal of Fungi*, 7 (2), 73.


Yang, Q., Chen, W.Y., Jiang, N., Tian, C.M. 2019. *Nectria*-related fungi causing dieback and canker diseases in China, with *Neothyronectria citri* sp. nov. described. *MycKeys*, 56, 49–66.

Yang, Q., Du, Z., Liang, Y.M., Tian, C.M. 2018. Molecular phylogeny of *Nectria* species associated with dieback and canker diseases in China, with a new species described. *Phytotaxa*, 356 (3), 199–214.


APPENDIX. PHOTO PLATES



A: An infected twig *Abies spectabilis* with diseased needles showing characteristic symptom; B: Colony of *Neonectria neomacrospora* cultured from infected part and reverse view of culture plate; C: Conidia of *Neonectria neomacrospora*.



A: An infected leaf showing characteristic symptoms of powdery mildew on *Alnus nepalensis*; B: Conidia of *Golovinomyces eichloracearum*.




A: An infected leaf of *Eucalyptus camaldulensis* showing characteristic symptoms; B: Colony of *Calonectria reteaudii* cultured from infected part and reverse view of culture plate; C: Conidia of *Calonectria reteaudii*.



A and B: Upper and lower surface of infected leaves of *Acer laevigatum* showing characteristic symptom; C: Colony of *Aureobasidium apocryptum* cultured from infected part and reverse view of culture plate; D: Conidia with conidiophores of *Aureobasidium apocryptum*.



A: An infected trunk of *Alnus nepalensis* showing bleeding symptom; B: Colony of *Phytophthora alni* cultured from infected part and reverse view of culture plate; C: Zoosporangium.



A and B: An infected shoot and needles of *Pinus roxburghii*; C: Colony of *Dothistroma septosporum* cultured from infected part and reverse view of culture plate; D: Conidia of *Dothistroma septosporum*.

